



		<ul style="list-style-type: none"> <li>• Solu-Cortef 100mg IM x 1</li> <li>• Diphenhydramine 50mg IM x 1</li> </ul>			
		<ul style="list-style-type: none"> <li>➤ Place on telemetry and obtain BP, HR, RR, &amp; SpO<sub>2</sub> <ul style="list-style-type: none"> <li>• <b>Time 0:</b> Infuse 0.5 mg/kg [XYZStudyDrug] IV in 100mL 0.9% NaCl over 40min (study physician will remain on site until end of infusion). Study team will order study drug, &amp; CRU nurse to pick up from the pharmacy.</li> <li>• VS q15min for duration of infusion</li> </ul> </li> </ul>			
		<ul style="list-style-type: none"> <li>➤ <b>Time +40min (end infusion):</b> <ul style="list-style-type: none"> <li>• Monitor BP, HR, RR, &amp; SPO<sub>2</sub> via telemetry. Notify study team for any S/S of adverse reaction immediately</li> <li>• VS q15min x 1hr</li> </ul> </li> </ul>			
		<ul style="list-style-type: none"> <li>➤ <b>Time +100min (1hr post-end of infusion):</b> <ul style="list-style-type: none"> <li>• D/C IV and telemetry and provide lunch if participant is not sedated</li> </ul> </li> </ul>			
		<ul style="list-style-type: none"> <li>➤ <b>Post-Infusion Procedures:</b> <ul style="list-style-type: none"> <li>• Notify PI for psychiatric emergencies (suicidality, psychosis, agitation)</li> <li>• Lorazepam 2mg po q4h prn agitation/anxiety</li> <li>• Lorazepam 2mg IM q4h prn agitation/anxiety</li> <li>• Haloperidol 5 mg po q2h prn psychosis</li> <li>• Haloperidol 5mg IM q2h prn psychosis <ul style="list-style-type: none"> <li>• Prior to giving Lorazepam or Haloperidol, notify Dr. [PI]</li> <li>• Notify physician who administered infusion for medical emergencies: SBP &gt;160, DBP &gt;110, or HR &gt;100</li> </ul> </li> </ul> </li> </ul>			
		<ul style="list-style-type: none"> <li>➤ <b>Time +120 min (2 hrs post-start of infusion):</b> <ul style="list-style-type: none"> <li>• Study team will evaluate participant and, if stable for discharge, will discharge home</li> </ul> </li> </ul>			